

## **Fluorescein-Labeled Antibody To Human IgA+IgG+IgM (H+L)**

### ***Produced in Goat***

**Catalog No.**  
**5230-0290 (02-10-17)**

**Size**  
**0.5 mg**

#### **DESCRIPTION**

This broad spectrum reagent contains a mixture of 3 antibodies that recognize: human IgA, IgG and IgM. Affinity purified antibody isolated from a pool of serum from goats immunized with purified human IgA, IgG, and IgM was labeled with fluorescein isothiocyanate.

#### **FORM/STORAGE**

Lyophilized. Store at 2-8°C until rehydrated. Stable for a minimum of 1 year when stored at 2-8°C.

#### **STABILIZER AND PRESERVATIVE**

Goat serum and/or bovine serum albumin (BSA) is added as a protein stabilizer. No preservative added. Additional biological protection may be provided with 0.1% sodium azide. Non-sterile.

#### **ANTIBODY CONCENTRATION**

The concentration of affinity purified antibody is 0.5 mg as determined by UV absorbance at 280 nm.

#### **F/P RATIO**

The fluorescein:antibody protein ratio is between 3:1 and 7:1.

#### **SPECIFICITY/CROSS REACTIVITY**

Tested by gel diffusion and ELISA techniques as applicable. This product reacts specifically with human IgA, IgG, and IgM and may recognize other immunoglobulin types that have light chains in common. Reactivity to IgG subclasses has not been tested. Antibodies to human IgA, IgG, and IgM may cross-react with immunoglobulins of other mammalian species if common binding sites are shared.

#### **REHYDRATION AND STORAGE**

Note: Rehydration of antibodies in TBS or buffers other than those listed here is not recommended.

**Rehydration:** Rehydrate with 1 mL of reagent quality water. Rotate the vial until the lyophilized pellet is totally dissolved.

**Storage:** This product may be stored for up to 1 week refrigerated; thereafter, it should be stored frozen. Stable for a minimum of 1 year at - 20 °C.

#### **SUGGESTED WORKING DILUTIONS**

Optimal working concentrations should be determined experimentally. Prepare working dilutions in TBS or other buffer such as 5X Detector Block or Milk/Diluent Blocking Solution (See RELATED PRODUCTS) immediately before use.

These buffers are not recommended for long term storage. A suggested starting dilution of 1:10-1:100 is recommended for most applications. In many cases, the antibody may be diluted further than indicated.

#### **PRODUCT SAFETY AND HANDLING**

This product is considered non-hazardous as defined by The Hazard Communication Standard (29 CFR 1910.1200). Avoid contact with skin and eyes. In case of contact or spillage, clean with copious amounts of water. Disposal via sanitary sewer.

#### **RELATED PRODUCTS**

#### **CAT. NO.**

KPL 5X Detector Block	5920-0004 (71-83-00)
KPL Milk Diluent/Blocking Solution	5140-0011 (50-82-01)
KPL Coating Solution	5150-0014 (50-84-00)
KPL Wash Solution	5150-0008 (50-63-00)
KPL 10% BSA Diluent/Block	5140-0006 (50-61-00)